

ASTROMETRY WITH THE *HUBBLE SPACE TELESCOPE*: A PARALLAX OF THE FUNDAMENTAL DISTANCE CALIBRATOR δ CEPHEI¹

G. FRITZ BENEDICT,² B. E. MCARTHUR,² L. W. FREDRICK,³ T. E. HARRISON,⁴ C. L. SLESNICK,³ J. RHEE,³ R. J. PATTERSON,³
M. F. SKRUTSKIE,³ O. G. FRANZ,⁵ L. H. WASSERMAN,⁵ W. H. JEFFERYS,⁶ E. NELAN,⁷ W. VAN ALTENA,⁸ P. J. SHELUS,²
P. D. HEMENWAY,⁹ R. L. DUNCOMBE,¹⁰ D. STORY,¹¹ A. L. WHIPPLE,¹¹ AND A. J. BRADLEY¹²

Received 2002 February 11; accepted 2002 May 28

ABSTRACT

We present an absolute parallax and relative proper motion for the fundamental distance scale calibrator δ Cep. We obtain these with astrometric data from FGS 3, a white-light interferometer on the *Hubble Space Telescope* (*HST*). Utilizing spectrophotometric estimates of the absolute parallaxes of our astrometric reference stars and constraining δ Cep and reference star HD 213307 to belong to the same association (Cep OB6), we find $\pi_{\text{abs}} = 3.66 \pm 0.15$ mas. The larger than typical astrometric residuals for the nearby astrometric reference star HD 213307 are found to satisfy Keplerian motion with $P = 1.07 \pm 0.02$ yr, a perturbation and period that could be due to an F0 V companion ~ 7 mas distant from and ~ 4 mag fainter than the primary. Spectral classifications and *VRIJHKT₂M* and DDO51 photometry of the astrometric reference frame surrounding δ Cep indicate that field extinction is high and variable along this line of sight. However the extinction suffered by the reference star nearest (in angular separation and distance) to δ Cep, HD 213307, is lower and nearly the same as for δ Cep. Correcting for color differences, we find $\langle A_V \rangle = 0.23 \pm 0.03$ for δ Cep and hence an absolute magnitude $M_V = -3.47 \pm 0.10$. Adopting an average V magnitude, $\langle V \rangle = 15.03 \pm 0.03$, for Cepheids with $\log P = 0.73$ in the large Magellanic Cloud (LMC) from Udalski et al., we find a V -band distance modulus for the LMC, $m - M = 18.50 \pm 0.13$, or 18.58 ± 0.15 , where the latter value results from a highly uncertain metallicity correction. These agree with our previous RR Lyr *HST* parallax-based determination of the distance modulus of the LMC.

Key words: astrometry — distance scale — Magellanic Clouds — stars: distances — stars: individual (δ Cephei)

1. INTRODUCTION

Many of the methods used to determine the distances to remote galaxies and ultimately the size, age, and shape of the universe itself depend on our knowledge of the distances to local objects. The most important of these are the Cepheid variable stars. Considerable effort has gone into determining the absolute magnitudes, M_V , of these objects (see the comprehensive review by Feast 1999). Given that the distances of all local Cepheids, except Polaris, are in

excess of 250 pc, most of these M_V determinations rely on large number statistics, for example, Groenewegen & Oudmaijer (2000), Lanoix, Paturel, & Garnier (1999), and Feast (1997). Gieren, Barnes, & Moffett used Cepheid surface brightness to estimate distances and absolute magnitudes. For Cepheid variables, these determinations are complicated by dependence on color and metallicity. Only recently have relatively high precision trigonometric parallaxes been available for a very few Cepheids (the prototype, δ Cep, and Polaris) from *Hipparcos* (δ Cep=HIP 110991; Perryman et al. 1997). We have determined the parallax of δ Cep with FGS 3 on the *Hubble Space Telescope* (*HST*) with significantly higher precision. Additionally, our extensive investigation of the astrometric reference stars provides an independent estimation of the line-of-sight extinction to δ Cep, a significant contributor to the uncertainty in its absolute magnitude, M_V .

In this paper we briefly discuss (§ 2) data acquisition, analysis, and an improved FGS 3 calibration, present the results of spectrophotometry of the astrometric reference stars required to correct our relative parallax to absolute (§ 3), and derive an absolute parallax for δ Cep (§ 4.4). Finally, we calculate an absolute magnitude for δ Cep (§ 5.3) and apply it to derive a distance modulus for the Large Magellanic Cloud (§ 5.4).

Bradley et al. (1991) and Nelan et al. 2001 provide an overview of the FGS 3 instrument and Benedict et al. (1999) describe the fringe-tracking mode astrometric capabilities of FGS 3, along with data acquisition and reduction strategies used in the present study. We time-tag our data with a modified Julian Date, $\text{mJD} = \text{JD} - 2,444,000.5$.

¹ Based on observations made with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

² McDonald Observatory, RLM 15.038, University of Texas, Austin, TX 78712.

³ Department of Astronomy, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903-0818.

⁴ Department of Astronomy, New Mexico State University, Box 30001, MSC 4500, Las Cruces, NM 88003-8001.

⁵ Lowell Observatory, 1400 West Mars Hill Road, Flagstaff, AZ 86001.

⁶ Department of Astronomy, RLM 15.038, University of Texas, Austin, TX 78712.

⁷ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218.

⁸ Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520.

⁹ Department of Oceanography, University of Rhode Island, Kingston, RI 02881.

¹⁰ Department of Aerospace Engineering, University of Texas, Austin, TX 78712.

¹¹ Aerospace Engineering Division, Jackson and Tull, Suite 200, 7375 Executive Place, Seabrook, MD 20706.

¹² Spacecraft System Engineering Services, P.O. Box 91, Annapolis Junction, MD 20706.

2. OBSERVATIONS AND DATA REDUCTION

Figure 1 shows the distribution in right ascension and declination of the five reference stars and δ Cep. Seven sets of data were acquired, spanning 2.44 yr, for a total of 127 measurements of δ Cep and reference stars. Each data set required approximately 40 minutes of spacecraft time. The data were reduced and calibrated as detailed in Benedict et al. (1999) and McArthur et al. (2001). In a recent paper (Benedict et al. 2002) we described a technique used for these data in which we employ a neutral density filter to relate astrometry of very bright targets to faint reference stars. At each epoch we measured reference stars and the target, δ Cep, multiple times. We do this to correct for intra-orbit drift of the type seen in the cross-filter calibration data in our recent paper reporting a parallax for RR Lyr (Benedict et al. 2002, Fig. 1). Data sets 2 and 5 were each also afflicted with an episode of nonmonotonic drift, possibly due to mechanism (filter wheel) motion. Fortunately, the strategy of multiple repeats of the observation sequence within each data set permitted generally satisfactory correction.

All these observations were acquired with FGS 3, and all before 1998. Since 1999, the prime astrometer aboard the *Hubble Space Telescope* (*HST*) has been FGS 1r, installed during the 1999 servicing mission. To calibrate the optical field angle distortion (OFAD) for FGS 1r we have, during the last three years, secured over 30 observation sets of M35, our calibration target field. These new observations have extended our time base to over 10 years and allowed us to refine the proper motions of calibration stars in M35. This has resulted in a more precise star catalog, which in turn has allowed us to improve our FGS 3 OFAD. Applying this revised calibration to the Barnard's star data presented in Benedict et al. (1999), we find a parallax difference of 0.07 mas and a proper-motion difference of 0.7 mas yr^{-1} , each a

change of about 1 part in 10,000. We use this new calibration for these δ Cep data.

3. SPECTROPHOTOMETRIC ABSOLUTE PARALLAXES OF THE ASTROMETRIC REFERENCE STARS

Because the parallax determined for δ Cep will be measured with respect to reference frame stars that have their own parallaxes, we must either apply a statistically derived correction from relative to absolute parallax (van Altena, Lee, & Hoffleit 1995) or, preferably, estimate the absolute parallaxes of the reference frame stars seen in Figure 1. With colors, spectral type, and luminosity class for a star one can estimate the absolute magnitude, M_V , and V -band absorption, A_V . The absolute parallax is then

$$\pi_{\text{abs}} = 10^{-(V-M_V+5-A_V)/5}. \quad (1)$$

The luminosity class is generally more difficult to determine than the spectral type (temperature class). However, the derived absolute magnitudes are critically dependent on the luminosity class. To confirm the luminosity classes we employ the technique used by Majewski et al. (2000) to discriminate between giants and dwarfs for stars later than $\sim G5$, an approach whose theoretical underpinnings are discussed by Paltoglou & Bell (1994).

3.1. Photometry

Our bandpasses for reference star photometry include *BVRI*, *JHK* (from preliminary 2MASS¹³ data), and Washington-DDO filters M, 51, and T₂ (obtained at McDonald Observatory with the 0.8 m prime focus camera). We transform the 2MASS *JHK* to the Bessell (1979) system by using the transformations provided in Carpenter (2001). In Tables 1 and 2 we list the visible, infrared, and Washington-DDO photometry for the δ Cep reference stars, DC-2 through DC-7. DC-2 was too bright for 2MASS and our Washington-DDO and *RI* photometric techniques.

3.2. Spectroscopy

The spectra from which we estimated spectral type and luminosity class come from the New Mexico State University Apache Point Observatory.¹⁴ For all but reference star DC-2 classifications were obtained by a combination of template matching and line ratios. Differing classifications for reference star DC-2 (HD 213307) have been reported in the literature. We consider the classifications of both Lutz & Lutz (1977) and Savage et al. (1985). Table 3 contains a list of the spectral types and luminosity classes for our reference stars, rank-ordered by estimated distance. We discuss our estimation of the $\langle A_V \rangle$ in the next subsection.

In Figure 2 we plot the Washington-DDO photometry along with a dividing line between dwarfs and giants (Paltoglou & Bell 1994). The boundary between giants and dwarfs is actually far fuzzier than suggested by the solid line and complicated by the photometric transition from dwarfs to giants through subgiants. This soft boundary is readily




FIG. 1.— δ Cep and astrometric reference stars. Symbol size is indicative of V magnitude (Table 1). The numbers within each identification box are the $\langle A_V \rangle$ per star from Table 4, § 3.3, and $E(B-V)$ per 100 pc from Table 3.

¹³ The Two Micron All-Sky Survey is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center at California Institute of Technology.

¹⁴ The Apache Point Observatory 3.5 m telescope is owned and operated by the Astrophysical Research Consortium.

TABLE 1
ASTROMETRIC REFERENCE STARS: PHOTOMETRY

ID	V	$B-V$	$V-R$	$V-I$	$V-K$
DC-2 ^a	6.30 ± 0.02	-0.04 ± 0.02
DC-3.....	13.47 ± 0.02	1.60 ± 0.10	1.10 ± 0.04	2.17 ± 0.04	4.22 ± 0.06
DC-4.....	12.68 ± 0.02	1.83 ± 0.10	1.34 ± 0.03	2.56 ± 0.03	4.93 ± 0.06
DC-5.....	13.68 ± 0.02	1.34 ± 0.10	0.96 ± 0.04	1.94 ± 0.04	3.69 ± 0.06
DC-7.....	14.18 ± 0.02	1.60 ± 0.10	1.11 ± 0.04	2.22 ± 0.04	4.19 ± 0.07

^a Lutz & Lutz 1977.

TABLE 2
ASTROMETRIC REFERENCE STARS: NEAR-IR AND WASHINGTON-DDO PHOTOMETRY

ID	K	$J-H$	$J-K$	$M-T_2$	$M-51$
DC-2.....	-0.17 ± 0.02	0.02 ± 0.02
DC-3.....	9.25 ± 0.02	0.76 ± 0.02	0.93 ± 0.03	2.22 ± 0.02	-0.07 ± 0.02
DC-4.....	7.75 ± 0.02	0.93 ± 0.02	1.21 ± 0.03	2.58 ± 0.01	-0.1 ± 0.01
DC-5.....	9.99 ± 0.02	0.71 ± 0.02	0.84 ± 0.03	1.97 ± 0.02	0.01 ± 0.03
DC-7.....	9.99 ± 0.02	0.67 ± 0.02	0.88 ± 0.03	2.25 ± 0.01	-0.05 ± 0.02

apparent in Majewski et al. (2000, Fig. 14). Objects just above the heavy line are statistically more likely to be giants than objects just below the line. Reference stars DC-2 and DC-5 have spectral types that are too early for this discriminant to work properly. The photometry is consistent with a giant or subgiant classification for the other reference stars.

3.3. *Interstellar Extinction*

To determine interstellar extinction we first plot these stars on several color-color diagrams. A comparison of the relationships between spectral type and intrinsic color against measured colors provides an estimate of reddening. Figure 3 contains $V-R$ vs. $V-K$ and $V-I$ vs. $V-K$ color-color diagrams and reddening vectors for $A_V = 1.0$. Also plotted are mappings between spectral type and luminosity class V and III from Bessell & Brett (1988) and Cox (2000, hereafter AQ00), again with reddening vectors and the loci of luminosity classes V and III stars. Figure 3, along with the estimated spectral types, provides measures of the reddening for each reference star.

TABLE 3
ASTROMETRIC REFERENCE STARS: SPECTRAL CLASSIFICATIONS AND SPECTROPHOTOMETRIC PARALLAXES

ID	Sp.	V	M_V	A_V	π_{abs} (mas)
DC-2.....	B7 IV ^a	6.30	-0.85 ± 0.4^b	0.28	4.2 ± 0.8
DC-2.....	B8 III ^c	6.30	-1.35 ± 0.4^b	0.09	3.2 ± 0.6
DC-2.....	B7-B8 III-V ^d	6.30	-1.10 ± 0.1	0.21	3.70 ± 0.26
DC-3.....	K1 III	13.47	0.6 ± 0.4	1.63	0.6 ± 0.1
DC-7.....	G8 III	14.18	0.8 ± 0.4	2.05	0.5 ± 0.1
DC-4.....	K3 III	12.68	0.3 ± 0.4	2.07	0.9 ± 0.2
DC-5.....	G1 III	13.68	0.9 ± 0.4	2.13	0.7 ± 0.1

^a Lutz & Lutz 1977.

^b Wegner 2000.

^c Savage et al. 1985.

^d From membership in Cep OB6 (de Zeeuw et al. 1999).

Assuming an $R = 3.1$ galactic reddening law (Savage & Mathis 1979), we derive A_V values by comparing the measured colors (Tables 1 and 2) with intrinsic $V-R$, $V-I$, $J-K$, and $V-K$ colors from Bessell & Brett (1988) and AQ00. Specifically, we estimate A_V from four different ratios, each derived from the Savage & Mathis (1979) reddening law: $A_V/E(V-R) = 4.83$; $A_V/E(V-K) = 1.05$; $A_V/E(J-K) = 5.80$; and $A_V/E(V-I) = 2.26$. These A_V are collected in Table 4. Colors and spectral types are




Fig. 2.— M -DDO51 ($M-51$) vs. $M-T_2$ color-color diagram for reference stars DC-3 through DC-7. The solid line is the division between luminosity class V and luminosity class III stars. Giants are above the line, dwarfs below. The reddening vector is for $A_V = 1.0$. The numbers are the reference star IDs plotted at the observed values. The circles are dereddened values, based on the $\langle A_V \rangle$ per star from Table 4.




FIG. 3.— $V-R$ vs. $V-K$ and $V-I$ vs. $V-K$ color-color diagrams for reference stars DC-3 through DC-7. The dashed line is the locus of dwarf (luminosity class V) stars of various spectral types; the dot-dashed line is for giants (luminosity class III). The reddening vector is for an $A_V = 1$.

inconsistent with a field-wide average $\langle A_V \rangle$ for the δ Cep field. The spatial distribution of the average reddening star to star is shown in Figure 1. A simple uniform extinction would predict a correlation between A_V and distance, with more distant objects having higher A_V . This correlation is absent in Table 3, suggesting that either the extinction or the distances are in error. Alternatively, a patchy distribution of the interstellar matter would destroy any correlation, a distinct possibility for this field at Galactic latitude $l = 0^\circ.5$. As we shall see, the reddening for reference star DC-2 (discussed in § 4.4), is of critical importance to an estimate of the reddening at the location of δ Cep.

3.4. Reference Frame Absolute Parallaxes

We have prior knowledge that reference star DC-2 is thought to be physically associated with δ Cep. Hoffleit &

TABLE 4
ASTROMETRIC REFERENCE STARS: A_V FROM SPECTROPHOTOMETRY

ID	A_V ($V-I$)	A_V ($V-R$)	A_V ($V-K$)	A_V ($J-K$)	$\langle A_V \rangle^a$
DC-3.....	1.83	1.43	1.80	1.45	1.63 ± 0.12
DC-7.....	2.35	1.98	2.13	1.74	2.05 ± 0.15
DC-4.....	2.06	1.90	2.03	2.32	2.07 ± 0.10
DC-5.....	2.37	2.02	1.96	2.15	2.13 ± 0.10
$\langle A_V \rangle^b$	2.15	1.83	1.98	1.91	1.97 ± 0.13

^a Average by star.

^b Average by color index.

Jaschek 1982 (hereafter BSC82) note common proper motion with δ Cep and that DC-2=ADS 15987C (δ Cep=ADS 15987A), while de Zeeuw et al. (1999) include both δ Cep and DC-2 in the newly discovered Cep OB6 association. Consequently, we first explored the astrometric properties of the four remaining reference stars, solving for parallax and proper motion of each in turn relative to a reference frame defined by the other three reference stars. We obtained no significant improvement in χ^2 by allowing any reference star (other than DC-2) to have a proper motion relative to the other three. In each case the spectrophotometric parallaxes discussed below entered the solution as observations.

We derive absolute parallaxes with M_V values from AQ00 and the $\langle A_V \rangle$ obtained from the photometry. These are listed in Table 3, with three possible values for reference star DC-2, two depending on past spectral classifications. The last value for DC-2 is derived by constraining it to belong to the Cep OB6 association (see § 4.4 below). The weighted average absolute parallax for the reference frame is $\langle \pi_{\text{abs}} \rangle = 0.77$ mas, including the highest-weight parallax determination for DC-2, and 0.63 mas without DC-2. Statistically, DC-2 has very little weight in our reference frame. Nonetheless, it is astrometrically critical, as discussed in (§ 4.4), below.

4. ABSOLUTE PARALLAX OF δ CEPHEI

4.1. Astrometric Model

With the positions measured by FGS 3 we determine the scale, rotation, and offset plate constants relative to an arbitrarily adopted constraint epoch (the so-called master plate) for each observation set (the data acquired at each epoch). The mJD of each observation set is listed in Table 5, along with the magnitude measured by the FGS 3 (zero point provided by Barnes et al. 1997), a phase (based on $P = 5.366316$ days), and a $B-V$ estimated by comparison with the UBV photometry of Barnes et al. (1997). The δ Cep reference frame contains five stars. We employ the six-parameter model discussed in McArthur et al. (2001) for those observations. For the δ Cep field four of the reference stars are significantly redder than the science target and one is bluer. Hence, we apply the corrections for lateral color discussed in Benedict et al. (1999).

As in all our previous astrometric analyses, we employ GAUSSFIT (Jefferys, Fitzpatrick, & McArthur 1987) to minimize χ^2 . The solved equations of condition for

TABLE 5
 δ CEPHEI LOG OF OBSERVATIONS

Data Set	mJD	Phase ^a	V^b	$B-V^c$
1.....	49,908.60462	0.870	3.967	0.63
2.....	49,942.7928	0.241	3.910	0.64
3.....	50,087.07986	0.128	3.775	0.55
4.....	50,104.13513	0.306	3.989	0.69
5.....	50,630.66157	0.423	4.103	0.77
6.....	50,668.40227	0.456	4.131	0.79
7.....	50,799.52478	0.891	3.862	0.57

^a Phase based on $P = 5.366316$ days, $T_0 = 43,673.644$ (mJD) (Barnes et al. 1997).

^b Differential FGS photometry; technique described in Benedict et al. 1998. Zero point from Barnes et al. 1997.

^c Estimated from phase and Barnes et al. 1997 photometry.

δ Cep are

$$x' = x + lcx(B-V) - \Delta XF_x, \quad (2)$$

$$y' = y + lcy(B-V) - \Delta XF_y, \quad (3)$$

$$\xi = Ax' + By' + C + R_x(x'^2 + y'^2) - \mu_x \Delta t - P_\alpha \pi_x, \quad (4)$$

$$\eta = -Bx' + Ay' + F + R_y(x'^2 + y'^2) - \mu_y \Delta t - P_\delta \pi_y, \quad (5)$$

where x and y are the measured coordinates from *HST*, lcx and lcy are the lateral color corrections from Benedict et al. (1999), and $B-V$ are the $B-V$ colors of each star, including the variable $B-V$ of δ Cep (Table 5). Here ΔXF_x and ΔXF_y are the cross-filter corrections in x and y , applied to the observations of δ Cep and reference star DC-2. δ Cep has a full range of $0.2 < B-V < 0.6$. For this analysis we linearly interpolate between the 1995 and 1998 cross-filter calibrations (see Table 1, Benedict et al. 2002) as a function of δ Cep color. A and B are scale and rotation plate constants, C and F are offsets, R_x and R_y are radial terms, μ_x and μ_y are proper motions, Δt is the epoch difference from the mean epoch, P_α and P_δ are parallax factors, and π_x and π_y are the parallaxes in right ascension and declination. We obtain the parallax factors from a JPL Earth-orbit predictor (Standish 1990), upgraded to version DE405. Orientation to the sky is obtained from ground-based astrometry (USNO-A2.0 catalog, Monet 1998) with uncertainties in the field orientation $\pm 0:05$.

4.2. Assessing Reference Frame Residuals

The optical field angle distortion calibration (McArthur et al. 1997) reduces as-built *HST* telescope and FGS 3 distortions with magnitude $\sim 1''$ to below 2 mas over much of the FGS 3 field of regard. From histograms of the astrometric residuals (Fig. 4) we conclude that we have obtained correction at the ~ 1.5 mas level. The resulting reference frame “catalog” in ξ and η standard coordinates (Table 6) was determined with $\langle \sigma_\xi \rangle = 0.3$ and $\langle \sigma_\eta \rangle = 0.3$ mas.

Noting that the residual histograms have larger dispersions than we typically achieve, we plotted the δ Cep reference frame residuals against a number of spacecraft, instrumental, and astronomical parameters to determine whether there might be unmodeled—but possibly correctable—systematic effects at the 1 mas level. The plots against residual included x and y position within the pickle, radial distance from the pickle center, reference star V magnitude




FIG. 4.—Histograms of x - and y -residuals obtained from modeling δ Cep and the astrometric reference stars with eqs. (4) and (5). Distributions are fitted with Gaussians whose σ -values are noted in the plots.

and $B-V$ color, and epoch of observation. Except for reference star DC-2 (=HD 213307) discussed below, we saw no obvious trends, other than an expected increase in positional uncertainty with reference star magnitude. The largest residuals are associated with observations of reference stars DC-5 and DC-7 made during the two orbits with anomalous drift, discussed in § 2.

4.3. New Companion for HD 213307?

BSC82 notes a possible very short period companion ($P < 1$ day) to reference star DC-2=HD 213307. Such a companion would be undetectable by the FGS, either directly (changes in fringe structure e.g., Franz et al. 1998) or indirectly (astrometric perturbation of the primary). Nonetheless, for DC-2 we found clear long-term and non-linear trends in the residuals with time. Because *HST* provides only relative proper motions, we do not expect full

TABLE 6
 δ CEPHEI AND REFERENCE STARS: ASTROMETRY

ID	ξ^a	η^a	μ_x^b	μ_y^b
δ Cep.....	45.2774 ± 0.0003	104.9195 ± 0.0003	0.0174 ± 0.0002	0.0050 ± 0.0002
DC-2.....	37.4174 ± 0.0003	64.9353 ± 0.0003	0.0215 ± 0.0003	0.0037 ± 0.0003
DC-3 ^c	0.0000 ± 0.0003	0.0000 ± 0.0004
DC-4.....	131.7161 ± 0.0002	116.1802 ± 0.0003
DC-5.....	88.9765 ± 0.0003	-27.4270 ± 0.0003
DC-6.....	167.0683 ± 0.0003	76.5763 ± 0.0003

^a ξ and η are positions in arcseconds relative to DC-3.

^b μ_x and μ_y are relative motions in arcseconds per year.

^c R.A. = $22^{\text{h}} 29^{\text{m}} 04.59^{\text{s}}$, decl. = $+58^{\circ} 47' 40''.7$, J2000.0, epoch mJD 50,104.1363.

agreement with the *Hipparcos* absolute proper motions. However, the agreement between *Hipparcos* and *HST* for δ Cep was within the errors, while that for DC-2 was not (Table 8). To assess the possibility that the residuals are caused by a perturbation due to a longer period unseen companion we employed the model from our past binary star work,

$$\xi = Ax' + By' + C + R_x(x'^2 + y'^2) - \mu_x \Delta t - P_\alpha \pi_x - \text{ORBIT}_x, \quad (6)$$

$$\eta = -Bx' + Ay' + F + R_y(x'^2 + y'^2) - \mu_y \Delta t - P_\delta \pi_y - \text{ORBIT}_y, \quad (7)$$

where ORBIT is a function of the traditional astrometric orbital elements (Benedict et al. 2001). Because of the small number of epochs and observations we were unable to treat HD 213307 as we did Wolf 1062A (Benedict et al. 2001) and simultaneously solve for all the terms in equations (6) and (7). To obtain a solution we constrained all but the parallax, proper-motion, and ORBIT parameters to values previously determined using equations (4) and (5). The resulting orbital elements in Table 7 should be taken as highly preliminary. Figure 5 shows the residuals from the solution provided by equations (6) and (7) and the fitted orbit. Introducing these parameters to the modeling process reduced the residual histogram dispersions seen in Figure 4 from $\sigma_x = 1.5$ to $\sigma_x = 1.4$ mas and from $\sigma_y = 1.4$ mas to $\sigma_y = 1.0$ mas.

With $M = 4 M_\odot$ adopted as the mass of the B7–B8 primary (AQ00), the perturbation orbit size, $\alpha = 2.0$ mas, and the period, $P = 1.06$ yr, yield a component B mass, $M_B \sim 1.6 M_\odot$. If the companion is on the main sequence, this F0 V star would be approximately 7 mas (1.9 AU) dis-

tant from and ~ 3.8 mag fainter than the primary, consistent with the absence of a previous detection. Radial velocity variations of the primary would have an amplitude $K1 \sim 15$ km s⁻¹, difficult to detect in a B7–B8 III–IV star with $v \sin i = 135$ km s⁻¹ (BSC82).

4.4. Absolute Parallax of δ Cephei and HD 213307

In a quasi-Bayesian approach the lateral color and cross-filter calibration values were entered into the model as observations with associated errors. The reference star spectrophotometric absolute parallaxes also were input as observations with associated errors, not as hardwired quantities known to infinite precision. This approach allows us to incorporate any measurements relevant to our investigation. These include the *Hipparcos* parallaxes and proper motions of δ Cep and HD 213307 (HIP 110988) with errors.

Even though DC-2 is a binary with a poorly determined orbit, we find that we must include this reference star in the solution. An ideal astrometric reference frame would sur-

TABLE 7
 HD 213307: ELEMENTS OF PERTURBATION ORBIT

Parameter	Value
α (mas).....	2.0 ± 0.2
P (days).....	390 ± 9
P (yr).....	1.07 ± 0.1
T_0	2002.3 ± 0.1
e	0.35 ± 0.09
i (deg).....	36 ± 14
Ω (deg).....	100 ± 8
ω (deg).....	259 ± 7




FIG. 5.—Reference star DC-2 (HD 213307) astrometric residuals from modeling with eqs. (6) and (7). Each observational epoch (denoted as Data Set in Table 5) is coded with a unique symbol. The emphasized symbols are plotted on the derived orbit at the epoch of observation. The orbit is preliminary and suggests a $M = 1.6 M_\odot$ companion.

round δ Cep. From Figure 1 it is clear that DC-2 is necessary to minimize extrapolation of the scale determined by the bulk of the offset reference frame. Without DC-2 the reference frame geometry is even less ideal. DC-2 also provides a constraint on the cross-filter calibration, because both DC-2 and δ Cep were observed with the neutral density filter. Following a suggestion from the referee, we constrained the difference in parallax between δ Cep and DC-2, using our prior knowledge of their proximity (see § 3.4). From de Zeeuw et al. (1999, Figure 25) we can estimate that the 1σ dispersion in Galactic longitude for the OB association thought to contain both δ Cep and DC-2 is 3° . One can therefore infer that the 1σ dispersion in distance in this group is $3^\circ \text{ rad}^{-1} \sim 5\%$. Hence, the 1σ dispersion in the parallax difference between two group members (e.g., DC-2 and δ Cep) is

$$\Delta\pi = 0.05(3.7)\sqrt{2} \text{ mas} = 0.26 \text{ mas} , \quad (8)$$

where we have here adopted the mean parallax of Cep OB6, $\langle\pi\rangle = 3.7 \text{ mas}$, from de Zeeuw et al. 1999. This estimated difference allows us to assign a higher statistical weight to reference star DC-2 than if we had adopted either past spectral typing (lines 1 and 2 in Table 3) or the *Hipparcos* value. The parallax difference between δ Cep and DC-2 becomes an observation with associated error fed to our model, an observation used to estimate the parallax difference between the two stars, while solving for the parallax of δ Cep.

We obtain for δ Cep an absolute parallax $\pi_{\text{abs}} = 3.66 \pm 0.15 \text{ mas}$, and for DC-2, $\pi_{\text{abs}} = 3.65 \pm 0.15$. Introducing the Cep OB6 parallax dispersion constraint and the *Hipparcos* parallax and proper-motion measurements with their associated errors allows us to obtain a statistically significant result from a reference frame with very poor geometry. Our final δ Cep parallax differs by $\sim 1\sigma_{\text{HIP}}$ and by $\sim 3\sigma_{\text{HST}}$ from that measured by *Hipparcos*, $\pi_{\text{abs}} = 3.32 \pm 0.58 \text{ mas}$. Nordgren et al. (2002) have used long-baseline interferometry to measure the angular diameter of δ Cep and other Cepheids. Their calibration of the Barnes-Evans (1976) relationship (surface brightness vs. color index) yields a distance to δ Cep of $272 \pm 6 \text{ pc}$ and a parallax $\pi_{\text{abs}} = 3.68 \pm 0.08 \text{ mas}$. We note that our δ Cep absolute parallax and that of Nordgren et al. agree within each other's errors.

In Figure 6 we compare δ Cep astrometric parallax results from *HST*, *Hipparcos*, and Allegheny Observatory (AO). We plot the AO redetermination reported in Gatewood, Kiewiet de Jonge, & Stephenson (1998), not the original result (Gatewood, de Jonge, & Stephenson 1993). Also plotted is the Nordgren et al. (2002) result. Parallax and proper-motion results from *HST*, *Hipparcos*, and AO are collected in Table 8. Parallax and proper-motion results for reference star DC-2 are also presented in Table 8, because it was measured by *Hipparcos*.

5. DISCUSSION AND SUMMARY

5.1. *HST* Parallax Accuracy

Our parallax precision, an indication of our internal random error is often less than 0.4 mas. To assess our accuracy, or external error, we must compare our parallaxes with results from independent measurements. Following Gatewood et al. (1998), we plot all parallaxes obtained by the *HST* Astrometry Science Team with FGS 3 against those obtained by *Hipparcos*. Data for these seven objects are col-




FIG. 6.—Absolute parallax determinations for δ Cep. We compare astrometric results from *HST*, *Hipparcos*, and a recent determination from Allegheny Observatory (AO; Gatewood, Kiewiet de Jonge, & Persinger 1998). Nordgren et al. (2002) have derived a parallax from a new calibration of the Barnes-Evans relation, denoted BE.

lected in Table 9 and shown in Figure 7. We have not included four Hyades stars whose parallaxes are considered preliminary (van Altena et al. 1997). The dashed line is a weighted regression that takes into account errors in both input data sets.

The regression indicates a 2.5σ scale difference between the *Hipparcos* and *HST* results. In addition, the fitted χ^2 suggests that either the *Hipparcos* or *HST* errors are overstated. Some but not all of this discrepancy can be explained by pointing out that our log-log plot artificially spreads out these data, which are effectively in two clumps, large and

TABLE 8
FINAL δ CEPHEI AND HD 213307 (DC-2) ABSOLUTE PARALLAX AND RELATIVE PROPER MOTION COMPARED WITH PREVIOUS RESULTS

Parameter	δ Cephei	HD 213307
<i>HST</i> study duration	2.44 y	...
Number of observation sets.....	7	...
Reference stars (V)	12.06	...
Reference stars ($B-V$)	1.3	...
Reference stars (V) excl. DC-2.....	13.50	...
Reference stars ($B-V$) excl. DC-2.....	1.6	...
<i>HST</i> absolute parallax (mas).....	3.66 ± 0.15	3.65 ± 0.15
<i>Hipparcos</i> absolute parallax (mas).....	3.32 ± 0.58	3.43 ± 0.64
AO absolute parallax (mas).....	2.8 ± 0.7	...
<i>HST</i> relative proper motion		
(mas yr ⁻¹).....	17.4 ± 0.7	21.8 ± 1.2
In position angle (deg).....	73 ± 3	80 ± 5
<i>Hipparcos</i> absolute proper		
motion (mas yr ⁻¹).....	16.9 ± 3.1	16.1 ± 2.7
In position angle (deg).....	79 ± 14	74 ± 12

TABLE 9
HST AND *Hipparcos* ABSOLUTE PARALLAXES

Object	<i>HST</i> (mas)	<i>Hip</i> (mas)	<i>HST</i> Reference
Prox Cen	769.7 ± 0.3	772.33 ± 2.42	Benedict et al. 1999
Barnard's star	545.5 ± 0.3	549.3 ± 1.58	Benedict et al. 1999
Feige 24.....	14.6 ± 0.4	13.44 ± 3.62	Benedict et al. 2000
Wolf 1062	98.0 ± 0.4	98.56 ± 2.66	Benedict et al. 2001
RR Lyr	3.60 ± 0.20	4.38 ± 0.59	Benedict et al. 2002
δ Cep.....	3.66 ± 0.15	3.32 ± 0.58	This paper
HD 213307.....	3.65 ± 0.15	3.43 ± 0.64	This paper

small parallaxes. Further exploring this issue, we have conducted a fully Bayesian analysis of the data in question, with an additional factor that represents the degree to which the presumed errors agree with the fit. We find that the input standard deviations of the *HST* and *Hipparcos* data points may have been overstated by a factor of ~ 1.5 . It is not possible to decide which errors, *HST* or those of the *Hipparcos*, or some combination of the two, are overstated. However, *Hipparcos* errors have been subjected to many tests confirming their validity. Hence, it is likely that *HST* errors are overstated.

Regarding the scale difference, we compare two hypotheses: the null hypothesis ($a = 0, b = 1$) and the complex hypothesis ($a \neq 0, b \neq 1$). We find that the Bayes factor

against the null hypothesis (that the straight line in Figure 7 is correctly described by $a = 0, b = 1$) and in favor of a more complex hypothesis is about 30, depending on the priors we put on a and b . These 30 : 1 odds provide only modest support for the complex hypothesis, i.e., that there is a systematic scale deviation between the *HST* and *Hipparcos* results. We point out that the amount of data is very small and that much of the scale difference (if real) depends on the two largest parallaxes, which have a great deal of leverage.

Measured proper motions provide another argument against the reality of this scale difference. Because it is desirable to reduce the impact of proper-motion errors on an *HST-Hipparcos* comparison, we consider only two of the objects in Table 9 (Proxima Cen and Barnard's star). They have proper-motion vector lengths exceeding 3800 mas yr^{-1} . *HST* yields proper motions relative to a local reference frame and *Hipparcos*-produced absolute proper motions. The dissimilar approaches can result in unequal proper motions for the same object. However, by comparing *HST* with *Hipparcos*, the average difference between these proper-motion vectors is -0.01% , indicating a negligible scale difference.

5.2. Lutz-Kelker Bias

When using a trigonometric parallax to estimate the absolute magnitude of a star, a correction should be made for the Lutz-Kelker (LK) bias (Lutz & Kelker 1973). Because of the galactic latitude and distance of δ Cep and the scale height of the stellar population of which it is a member, we use a uniform space density for calculating the LK bias. An LK algorithm modified by Hanson (1979) that includes the power law of the parent population is used. A correction of $-0.015 \pm 0.01 \text{ mag}$ is derived for the LKH bias correction to the *HST* parallax of δ Cep. The LKH bias is small because $\sigma_{\pi}/\pi \sim 4\%$ is small.

5.3. Absolute Magnitude of δ Cephei

To obtain the intrinsic absolute magnitude of δ Cep we require the intensity-averaged apparent magnitude, the absolute parallax, and an estimate of interstellar extinction. To estimate the extinction we turn to reference star DC-2 (HD 213307). For this star we obtain an absolute parallax, $\pi_{\text{abs}} = 3.65 \text{ mas}$, between the two estimates, depending on past spectral type and luminosity class determinations (Table 3). From SIMBAD we collected $B-V$ for all B8 III and B7 IV stars with $V \leq 4$ (presumably the nearest and least reddened members of their respective classes). The average for B8 III was $\langle B-V \rangle = -0.09$ and for B7 IV $\langle B-V \rangle = -0.13$. We adopt an intermediate $(B-V)_0 = -0.11$. From the measured color, $B-V = -0.04$, we obtain a color excess $E(B-V) = 0.07$. Because a blue star can yield a different ratio, R , of total to selective absorption than a redder star, depending on the amount of color excess, we employ the formulation

$$R = 3.07 + 0.28(B-V)_0 + 0.04E(B-V) \quad (9)$$

and obtain for DC-2 $\langle A_V \rangle = 0.21 \pm 0.03$ (Laney & Stobie 1993). The absolute magnitude of HD 213307 is then $M_V = -1.10$. Given that the distances to δ Cep and DC-2 are the same within the errors, we adopt the same color excess, $E(B-V) = 0.07$ for δ Cep, yielding $(B-V)_0 = 0.59$ and $R = 3.24$. Therefore, $\langle A_V \rangle = 0.23 \pm 0.03$ for δ Cep. Our determination agrees with $\langle A_V \rangle = 0.25 \pm 0.06$ from




FIG. 7.—*Bottom*, *HST* absolute parallax determinations compared with *Hipparcos* for the seven targets listed in Table 9; *top*, *Hipparcos* residuals to the error-weighted impartial regression line. The error bars on the residuals are *Hipparcos* 1σ errors.

TABLE 10
 δ CEPHEI INTENSITY-WEIGHTED MAGNITUDES

Magnitude	J ^a	CK ^b	DDO ^c
$\langle V \rangle$	3.95	3.95	3.95
$\langle I \rangle$	2.99	3.18	...
$\langle V \rangle - \langle I \rangle$	0.96	0.75	...
A_V	0.23 ^d	...	0.25
A_I	0.14 ^d
$\langle V_0 \rangle$	3.73	3.73	3.70
$\langle I_0 \rangle$	2.84	3.06	...
M_V	-3.47 ± 0.10	-3.47 ± 0.10	...
M_I	-4.34 ± 0.10	-4.14 ± 0.10	...

^a Johnson V/I photometry from Moffett & Barnes 1985.

^b Transformation per Bessel 1979.

^c From <http://ddo.astro.utoronto.ca/cepheids.html>.

^d Estimated from § 5.3 and Savage & Mathis 1979 reddening curve.

Fernie et al. (1995).¹⁵ Because differential reddening effects are smaller at the I band, we use a Savage & Mathis $R = 3.1$ reddening law to provide $A_I = 0.14$.

We next require the intensity-averaged magnitude of δ Cep. From Feast & Catchpole (1997) we obtain an intensity-weighted average $\langle V \rangle = 3.954$. From the photometry of Moffett & Barnes (1985), we adopt a Johnson I -band intensity-averaged magnitude, $\langle I \rangle = 2.993$. With the *HST* absolute parallax from § 4.4 and $\langle A_V \rangle = 0.23 \pm 0.03$, we derive $M_V = -3.47 \pm 0.10$, including the LKH bias and associated uncertainty. Similarly, $M_{I(CK)} = -4.14 \pm 0.10$, where we have transformed the Johnson to the Cousins-Kron photometric system as per Bessell (1979). The intensity-averaged photometry and absolute magnitudes are collected in Table 10. Using surface brightness techniques, Gieren et al. obtain for δ Cep an absolute visual magnitude $M_V = -3.63 \pm 0.13$ and, for $\log P = 0.73$, $M_V = -3.55 \pm 0.11$ from their period-luminosity (P-L) calibration for 100 Cepheids. Using a large number of relatively imprecise *Hipparcos* parallaxes, Feast & Catchpole (1997) and Lanoix et al. (1999) derive slightly different P-L calibrations for Cepheid variable stars. For $\log P = 0.73$, Feast obtains $M_V = -3.48$, and Lanoix et al., $M_V = -3.46$. All determinations agree with our value within the errors.

5.4. Distance Modulus of the LMC

Uncertainty in the distance to the Large Magellanic Cloud (LMC) contributes a substantial fraction of the uncertainty in the Hubble constant (Mould et al. 2000). The *HST* Key Project on the Extragalactic Distance Scale (Freedman et al. 2001; Mould et al. 2000) and the Type Ia Supernovae Calibration Team (Saha et al. 1999) have adopted the distance modulus value $m - M = 18.5$. Values from 18.1 to 18.8 are reported in the current literature, with those less than 18.5 supporting the short distance scale and those greater than 18.5, the long distance scale. Comprehensive reviews of the methods can be found in Carretta et al. (2000), Gibson (1999), and Cole (1998).

Ideally, our absolute magnitude values for δ Cep, $M_V = -3.47 \pm 0.10$ and $M_{I(CK)} = -4.14 \pm 0.10$, combined with apparent magnitudes for LMC Cepheids (corrected for

reddening internal to the LMC) at $\log P = 0.73$ would yield a unique LMC distance modulus. However, there are several complications that render such a distance modulus suspect. The first of these is cosmic dispersion in Cepheid properties due to the finite width of the instability strip. As shown in Figure 5.13 of Binney & Merrifield (1998), lines of constant period are not horizontal in the H-R diagram. This complication motivates the determination of period-luminosity-color (P-L-C) relationships. Udalski et al. (1999) find a dispersion $\sigma = 0.07$ mag about the P-L-C relation for LMC Cepheids. The scatter about any P-L relation decreases markedly as one utilizes progressively redder bandpasses, which are less affected by internal reddening (Udalski et al. 1999 and Tanvir 1999). A reddening-free magnitude, W , yields the P-L relationship with the least dispersion; Udalski et al. (1999) find $\sigma = 0.08$ mag. Assuming that the remaining scatter in a P-L-C or reddening-free P-L relationship is cosmic scatter, we adopt 0.075 mag as the cosmic dispersion in the intrinsic properties of Cepheid variables, including the defining member of the class, δ Cep.

Varying amounts of line blanketing due to intrinsic metallicity differences can also affect the apparent magnitude of a star. A comprehensive discussion of its effect on the Cepheid P-L relation is presented by Freedman et al. (2001). We adopt their correction, used for both V and I band and differential with respect to the LMC, of -0.2 ± 0.2 mag dex^{-1} . Specific to our calibration of the LMC P-L relation, for δ Cep Andrievsky et al. (2002) find $[O/H] = +0.01$. The LMC has $[O/H] \sim -0.4$ (Kennicutt et al. 1998). Hence, if δ Cep had LMC metallicity, it would be 0.08 ± 0.08 mag brighter in V and I .

We adopt the OGLE LMC V and I -P-L relations (Udalski et al. 1999) because they are based on a very large number of Cepheids. Utilizing these to generate apparent absorption-corrected Cepheid magnitudes at $\log P = 0.73$, we obtain LMC V -band distance moduli $m - M = 18.50 \pm 0.13$ or, corrected for metallicity, $m - M = 18.58 \pm 0.15$. The corresponding I -band moduli are $m - M = 18.53 \pm 0.13$ and $m - M = 18.61 \pm 0.15$, respectively. The errors include a 0.075 mag rss (root sum squares) cosmic dispersion. We list these distance moduli in Table 11. Distance moduli corrected for metallicity are listed in the rightmost column as $m - M = f(Z)$. At this stage in its maturity, this term introduces as much uncertainty as correction.

All estimates agree within their respective errors with our recent determinations from *HST* astrometry of RR Lyr (Benedict et al. 2002). There we obtained values of 18.53 ± 0.10 and 18.38 ± 0.10 , where the range comes from differing $\langle V_0(\text{RR}) \rangle$ values for LMC RR Lyr variables found in the literature. We note that the errors associated with the Benedict et al. (2001) LMC distance moduli derived from RR Lyr neglected cosmic dispersion. According to Popowski & Gould (1998) this amounts to ~ 0.14 mag, increasing our distance moduli errors to 0.18 mag. Table 12 and Figure 8 summarize LMC distance modulus determinations based on *HST* astrometry and compare them with weighted averages of a number of results based on RR Lyr and Cepheid variables. A more comprehensive version of this figure, along with the corresponding complete table and references, summarizing over 80 determinations based on 21 independent methods, can be found on the Web.¹⁶ All

¹⁵ See <http://ddo.astro.utoronto.ca/cepheids.html>.

¹⁶ See <http://clyde.as.utexas.edu/SpAstNEW/head.ps>.

TABLE 11
LMC DISTANCE MODULI FROM THE OGLE^a CEPHEID PERIOD-LUMINOSITY RELATION

Wavelength	a	b	m	$m - M^b$	$m - M = f(Z)^{b,c}$
V	-2.760 ± 0.031	17.04 ± 0.021	15.03 ± 0.03	18.50 ± 0.13	18.58 ± 0.15
I	-2.962 ± 0.021	16.56 ± 0.014	14.40 ± 0.02	18.53 ± 0.13	18.61 ± 0.15

^a Udalski et al. 1999; $m = a \log P + b$.

^b Errors are rss errors in m from P-L, absolute magnitude (M) error, and 0.075 magnitude cosmic dispersion.

^c Freedman et al. 2001 metallicity correction, $\Delta(m - M) = -0.2 \pm 0.2 \text{ mag dex}^{-1}$.

LMC distance moduli based on *HST* astrometry are consistent with the LMC value $m - M = 18.50 \pm 0.10$ adopted by the *HST* Distance Scale Key Project (Freedman et al. 2001).

5.5. Summary

HST astrometry yields an absolute trigonometric parallax for δ Cep, $\pi_{\text{abs}} = 3.66 \pm 0.15$ mas. This high-precision result requires an extremely small Lutz-Kelker bias correction, -0.015 ± 0.01 mag. To reduce our astrometric residuals to near-typical levels requires that we model reference star DC-2 as a binary and constrain it and δ Cep to belong to the same stellar group, Cep OB6. Our astrometric results for DC-2 yield an extinction for that star of $\langle A_V \rangle = 0.21 \pm 0.03$. Correcting for color-dependent differences in R , we find $\langle A_V \rangle = 0.23 \pm 0.03$ for δ Cep. The dominant contributor to the error in the resulting absolute magnitude for δ Cep, $M_V = -3.47 \pm 0.10$, remains the parallax. We find an LMC V -band distance modulus $m - M = 18.50 \pm 0.13$, uncorrected for metallicity. This value is in agreement with our previous determinations with *HST* astrometry of RR Lyr and the value adopted by the *HST* Distance Scale Key Project (Freedman et al. 2001).

Support for this work was provided by NASA through grants GTO NAG5-1603 from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These results are based partially on observations obtained with the Apache Point Observatory

TABLE 12
 V -BAND LMC DISTANCE MODULI

No.	Object	Source	$m - M$
1.....	δ Cep	This paper, $m - M = f(Z)$	18.58 ± 0.15^a
2.....	RR Lyrae	Benedict et al. 2002	18.53 ± 0.18^b
3.....	All Cepheids	Benedict et al. 2002, Table 10	18.53 ± 0.07^c
4.....	δ Cep	This paper, $m - M \neq f(Z)$	18.50 ± 0.13^d
5.....	All RR Lyr	Benedict et al. 2002, Table 10	18.45 ± 0.08^e
6.....	RR Lyrae	Benedict et al. 2002	18.38 ± 0.18^f

^a LMC Cepheid $\langle V \rangle$ from Udalski et al. 1999; Freedman et al. 2001 metallicity correction.

^b Carretta et al. 2000 LMC $\langle V_{\text{RR}}(0) \rangle = 19.14$ (includes $+0.03$, $\langle [\text{Fe}/\text{H}] \rangle$ correction).

^c Weighted average. Error is standard deviation of the mean from four independent techniques.

^d LMC Cepheid $\langle V \rangle$ from Udalski et al. 1999; no metallicity correction.

^e Weighted average. Error is standard deviation of the mean from five independent techniques.

^f Udalski et al. 1999; LMC $\langle V_{\text{RR}}(0) \rangle = 18.94$ (with $+0.05$, the $\langle [\text{Fe}/\text{H}] \rangle$ correction, 18.99).

3.5 m telescope, which is owned and operated by the Astrophysical Research Consortium. This publication makes use of data products from the Two Micron All-Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center (IPAC) at California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France; the NASA/IPAC Extragalactic Database, which is operated by the Jet Propulsion Laboratory (JPL), California Institute of Technology, under contract with the National Aeronautics and Space Administration; and NASA's Astrophysics Data System Abstract Service. Thanks to Tom Barnes for helpful discussions and an early review of the text. Thanks to the many people in Danbury, CT, who have surfed through many waves of change (once Perkin-Elmer, then Hughes Aerospace, then Raytheon, now Goodrich) and continue to support the FGS, especially Linda Abramowicz-Reed. Finally, we thank Andy Gould for his careful and critical refereeing, and for his suggestion to relate the reference star DC-2 and δ Cep parallaxes to each other through their membership in Cep OB6.




Fig. 8.—Recent determinations of the V -band distance modulus of the Large Magellanic Cloud (see Table 12). Values labeled 3 and 5 are weighted averages of a number of independent determinations using Cepheids (3) and RR Lyr variables (5).

REFERENCES

- Andrievsky, S. M., et al. 2002, *A&A*, 381, 32
 Barnes, T. G., & Evans, D. S. 1976, *MNRAS*, 174, 489
 Barnes, T. G., Fernley, J. A., Frueh, M. L., Navas, J. G., Moffett, T. J., & Skillen, I. 1997, *PASP*, 109, 645
 Benedict, G. F., et al. 1999, *AJ*, 118, 1086
 ———. 2001, *AJ*, 121, 1607
 ———. 2000, *AJ*, 119, 2382
 ———. 2002, *AJ*, 123, 473
 ———. 1998, *AJ*, 116, 429
 Bessell, M. S. 1979, *PASP*, 91, 589
 Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
 Binney, J., & Merrifield, M. 1998, *Galactic Astronomy* (Princeton: Princeton Univ. Press)
 Bradley, A., Abramowicz-Reed, L., Story, D., Benedict, G., & Jefferys, W. 1991, *PASP*, 103, 317
 Carpenter, J. M. 2001, *AJ*, 121, 2851
 Carretta, E., Gratton, R. G., Clementini, G., & Fusi Pecci, F. 2000, *ApJ*, 533, 215
 Cole, A. A. 1998, *ApJ*, 500, L137
 Cox, A. N. 2000, *Allen's Astrophysical Quantities*, ed. Arthur N. Cox (4th ed.; New York: AIP) (AQ00)
 de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H. J., Brown, A. G. A., & Blaauw, A. 1999, *AJ*, 117, 354
 Feast, M. 1999, *PASP*, 111, 775
 Feast, M. W. 1997, *MNRAS*, 284, 761
 Feast, M. W., & Catchpole, R. M. 1997, *MNRAS*, 286, L1
 Fernie, J. D., Evans, N. R., Beattie, B., & Seager, S. 1995, *Inf. Bull. Variable Stars*, 4148, 1
 Franz, O. G., et al. 1998, *AJ*, 116, 1432
 Freedman, W. L., et al. 2001, *ApJ*, 553, 47
 Gatewood, G., de Jonge, K. J., & Stephenson, B. 1993, *PASP*, 105, 1101
 Gatewood, G., Kiewiet de Jonge, J., & Persinger, T. 1998, *AJ*, 116, 1501
 Gibson, B. K. 1999, preprint (astro-ph/9910547)
 Gieren, W. P., Barnes, T. G., & Moffett, T. J. 1993, *ApJ*, 418, 135
 Groenewegen, M. A. T., & Oudmaijer, R. D. 2000, *A&A*, 356, 849
 Hanson, R. B. 1979, *MNRAS*, 186, 875
 Hoffleit, D., & Jaschek, C. 1982, *The Bright Star Catalogue* (4th ed.; New Haven: Yale Univ. Obs.) (BSC82)
 Jefferys, W., Fitzpatrick, J., & McArthur, B. 1987, *Celest. Mech.* 41, 39
 Kennicutt, R. C., et al. 1998, *ApJ*, 498, 181
 Laney, C. D., & Stobie, R. S. 1993, *MNRAS*, 263, 921
 Lanoix, P., Paturel, G., & Garnier, R. 1999, *MNRAS*, 308, 969
 Lutz, T. E., & Lutz, J. H. 1977, *AJ*, 82, 431
 Lutz, T. E., & Kelker, D. H. 1973, *PASP*, 85, 573
 Majewski, S. R., Ostheimer, J. C., Kunkel, W. E., & Patterson, R. J. 2000, *AJ*, 120, 2550
 McArthur, B., Benedict, G. F., Jefferys, W. H., & Nelan, E. 1997, in *Proc. 1997 HST Calibration Workshop*, ed. S. Casertano, R. Jędrzejewski, T. Keyes, & M. Stevens (Baltimore: STScI)
 McArthur, B. E., et al. 2001, *ApJ*, 560, 907
 Moffett, T. J., & Barnes, T. G. 1985, *ApJS*, 58, 843
 Monet, D. G. 1998, *BAAS* 193(112.003)
 Mould, J. R., et al. 2000, *ApJ*, 529, 786
 Nelan, E. 2001, *Fine Guidance Sensor Instrument Handbook*, Version 10 (Baltimore: STScI)
 Nordgren, T. E., Lane, B. F., Hindsley, R. B., & Kervella, P. 2002, *AJ*, 123, 3380
 Paltoglou, G., & Bell, R. A. 1994, *MNRAS*, 268, 793
 Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
 Popowski, P., & Gould, A. 1998, *ApJ*, 506, 259
 Saha, A., Sandage, A., Tammann, G. A., Labhardt, L., Macchetto, F. D., & Panagia, N. 1999, *ApJ*, 522, 802
 Savage, B. D., Massa, D., Meade, M., & Wesselius, P. R. 1985, *ApJS*, 59, 397
 Savage, B. D., & Mathis, J. S. 1979, *ARA&A*, 17, 73
 Standish, E. M., Jr. 1990, *A&A*, 233, 252
 Tanvir, N. R. 1999, in *ASP Conf. Ser. 167: Harmonizing Cosmic Distance Scales in a Post-Hipparcos Era*, ed. D. Egret & A. Heck (San Francisco: ASP), 84
 Udalski, A., Szymanski, M., Kubiak, M., Pietrzynski, G., Soszynski, I., Wozniak, P., & Zebrun, K. 1999, *Acta Astron.*, 49, 201
 van Altena, W. F., Lee, J. T., & Hoffleit, E. D. 1995, *Yale Parallax Catalog* (4th ed.; New Haven: Yale Univ. Obs.)
 van Altena, W. F., et al. 1997, *ApJ*, 486, L123
 Wegner, W. 2000, *MNRAS*, 319, 771